

Evaluation of drug prescribing patterns and cost analysis of Anti-diabetic therapy in Bargarh district.

Proposed by DR. Saurabh Kumar Banerjee,

DEAN, SCHOOL OF PHARMACEUTICAL MANAGEMENT,

SUBMITTED BY Subham Kumar Meher

District Name: Bargarh, Odisha

Demographics

AREA (IN SQ. KM)	5,837
TOTAL POPULATION	1,481,255
POPULATION DENSITY (PER SQ. KM.)	254
PROPORTION OF RURAL POPULATION TO TOTAL POPULATION	89.87
PROPORTION OF URBAN POPULATION TO TOTAL POPULATION	10.13
SEX RATIO(PER 1000)	977
LITERACY RATE	76%

INTRODUCTION

- Diabetes is a condition which is a metabolic disorder that is characterised along with different co morbidities and requires a lifelong treatment and maintenance of levels of glucose.
- Regardless of the Government attempt to prevent inexcusable price mounting of drugs by incorporating National List of Essential Medicine (NLEM) and by enacting the Drug Price Control Order (DPCO), huge discrepancy in cost of drugs still occur .
- The pattern of prescription still is used as an important tool to understands the trends in the medication prescription and also provides an idea about the drug usage and the rational use of drugs .

OBJECTIVES OF THE STUDY

1. To study the drug prescribing pattern .
2. To study the different class of major pharmaceutical drugs prescribe for diabetes and comorbidities .
3. To access the demographic and about the patient.
4. To know usage of the anti-diabetic medications and rational use of it.
5. To know the cost involved in the therapy

Research Questions

- What is Drug prescribing pattern among patients?
- What is the cost involved in the antidiabetic therapy ?

RESEARCH METHODOLOGY

RESEARCH TYPE	Primary and secondary
STUDY DESIGN	Descriptive and Analytical
SAMPLING METHOD	Random sampling
STUDY AREA	Bargarh district , Odisha
SAMPLE SIZE	110
DATA COLLECTION	Quantitative data
RESEARCH INSTRUMENT (TOOLS AND TECHNIQUES)	MS Excel
STUDY DURATION	2 months

Methods and Materials.


1. This study is conducted by collecting prescription from 110 participants and the sample are selected based on random sampling from different parts of Bargarh district and some of the prescriptions are collected from the different hospitals in the regions.
2. Permission from the hospital was taken prior to this study.
3. The sample size of the study is 110

Parameters of study.

- Categorisation of type of diabetes.
- Average anti-diabetic drugs prescribed per prescription.
- Which is the commonest and most frequent anti-diabetic drugs prescribed.
- Cost involved in the medication.
- Prescription is written in either generic or branded form.
- All other medication which are used for comorbidities along with anti-diabetic therapy.


Results and Discussion.

- **Categorisation of diabetes.**




As the graph suggest Type 2 diabetes is the common form of diabetes and accounts for 93 % and there is the less significance of type 2 diabetes which accounts for 7%.

Distribution as per number of anti-diabetics drugs in the prescription




- The usage of one or more anti-diabetic drugs are used .which suggest the usage of combination drugs therapy in comparison to the mono therapy used in diabetes .

Distribution based on Gender.



Among the 110 participants 65% male have diabetic conditions and 35% female ie 38 have diabeteic condition in the study sample.

Distribution based on Age.




Among the majority of participants the prevalence of diabetes is more in the age group 41-60 i.e 49% and the least prevalence of diabetes is in the age group 20-40.

- **Family history of diabetes patients**


- As we know genetic factor play a major role in diabetes so number of patients who have family history of diabetes is 37% as comparison to person who do not have family history of diabetes i.e 52% and 11 % people don't know about their family history of diabetes.

HISTORY OF DM	PARTICIPANTS	PERCENTAGE
Yes	41	37%
No	57	52%
Not significant	12	11%


- Among the different class of oral hypoglycemic agents Sulphonylureas are most commonly used .
- We can see that biaguanides and sulphonylureas are used in more prevalently i.e 37% and 30% followed by biaguanides and then the FDC of sulphonyl ureas and biaguanides i.e 20%.



- Metformin is the most commonly prescribed drug molecule in antidiabetic therapy i.e 37%
- After that glimepiride is prescribed more frequently and the combination of both two glimepiride and metformin is the drug of choice in FDC .




- Insulin preparation accounted for 45% .
- The most frequently used insulin is short acting insulin .




Branded vs Generic drugs.

- The percentage of branded drugs prescribed under this study is around 96% and the generic drugs were only prescribed in 4% Which is very less.



Prescribing pattern of drugs of comorbid conditions

- The commonly prescribed medications for comorbid conditions is Enalpril which have a 48% prescribing pattern followed by aspirin and then amlodipine which is 16% and 12% respectively .



- **Health economics of diabetes**

- The spending on health in India by the government is less as compare to the different developed countries. In India private health care is predominant and government set up provide around 20%of care in India.
- In India the out of pocket expenditure for diabetes is catastrophic and is accounted by 10% household consumption in 45% Patients and 23% takes loan to meet this out of pocket expenditure.
- The cost of therapy increased due to the different comorbidities and the use of insulin and the private health care As also we can see the increase amount of use of insulin in the Bargarh district of Odisha which accounts for 45% which add a lot of burden in the people's monetary expenditure.

- There is a significant increase in the price of antidiabetic therapy .

Years	Price Hike(rupees)
Less than 5 years	5.44 to 7.45 per day
Between 5 to 10 years	9.5 to 24 .23per day
Between 10 to 15 years	12.69 to 52.12
More than 15 years	17.50 to 63-78

- The direct expenditure of diabetes medication increased by 37% between 1998 and 2005 and the mean duration of diabetes was estimated to be 10.4 years and on comparing the direct population direct cost has increased upto 266%

Discussion

- The results obtained the cases and the number of patients of type 2 diabetes is more in comparison to the type 1 diabetes and the increasing percentage of type 2 diabetes is alarming situation in Bargarh district.
- The percentage of male diabetic condition is more than female diabetic condition i.e 65% and 35% which can be attributed to the lifestyle
- The prevalence of diabetes increase between 41-60 which suggest they develop diabetes in the most productive year due to life style modifications and stress.

- The study suggest the strong domination of OHA over the insulin therapy but as we can see Insulin usage is 45% which is higher in comparison to the previous study in India.
- This suggest a change in drug prescribing pattern and shifting towards the Insulin therapy in comparison to the OHA therapy.
- A large percentage of drugs are prescribed in brand name i.e96% Which suggest the high awareness of brand name within physician and the use of generic drugs are less which add in the increase in cost burden of the patient.

- Dominant use of sulphonylureas and biaguanides as still the class of drug for choice which accounts around 30% and 37% respectively and the combination of both the classes of drugs are still used frequently .
- This suggest that sulphonylurea and biguanides are still the choice of treatment with in the physician.
- Despite of the use of sulphonylureas metformin(biaguanide) is the most common used drugs by the prescriber its precribe frequency is 37% which is higher than the other molecules of the drugs in the study.

- The above suggest the use of enalapril and aspirin followed by atorvastatin and amlodipine, which suggest that most of the patients is on polypharmacy so they should be educate well and the interactions between the different class of drugs should be avoided for the rational use of drugs.

Limitation of study

- It is a pilot study which is conducted in the district of Bargarh by taking different prescriptions and interviewing the patients about their diabetes condition.
- It is conducted in a duration of 2 months in 110 diabetic patients.

CONCLUSION

- The present study analyse the drug prescribing pattern in the Bargarh district and the different cost that are involved in the maintance of the diabetes .
- Oral hypoglycemic agents are still dominate but the usage of insulin therapy increased over the period of time which suggest the change in shift in the prescribing pattern .
- It also study the prescription pattern of the patient with or without comorbities in the diabetes condition and the rational usage of the drugs

- The study suggest the use of polypharmacy which is due to the different kind of comorbidities condition in the diabetes.
- To avoid the irrational use of drugs the other drugs usage should be minimised .
- Due to the polypharmacy and the use of branded generics in the prescription it leads to the increase in the cost of prescription and leads to the out of pocket expenditure of the patient which should be avoided to lessen the out of pocket expenditure of medication in the patients.

REFERENCES

- American Diabetes Association. Standards of medical care in diabetes. Diabetes Care. 2018;41(Suppl 1):S13-S27.
-] American Diabetes Association. Economic Costs of Diabetes in US 2002. Diabetes Care. 2003;26:917-32.
 - American Diabetes Association. Economic Costs of Diabetes in US 2012. Diabetes Care. 2013;36:1033-46
- Meher M, Srivastav A, Jiwane D, Keer A. A Cost analysis study of oral hypoglycaemic drugs available in Indian market. World J Pharm Pharm Sci. 2018;7(3):1237-45.
- Bhanu P, Prasanand S, Divyashanthi CM, Annabelle R, Pandiamunian J. Prescribing pattern and WHO core prescribing indicators in post-operative patients of gynaecology department of tertiary care teaching hospital. Int J Basic Clin Pharmacology.
- Ahmed AM. History of diabetes mellitus. *Saudi Med J* 2002.
- Dipiro JT, Talbert RL, Gary CY, Matzke GR, Wells BG, Michael pharmacotherapy: A pathophysiologic approach
- Prescribing pattern and efficacy of anti-diabetic drugs in maintaining optimal glycemic levels in diabetic patients.
- American Diabetes Association. Diagnosis and classification of Diabetes mellitus. Diabetes care.

- Dipiro JT, Talbert RL, Gary CY, Matzke GR, Wells BG, Michael P. Pharmacotherapy: A pathophysiologic approach.
- Kastor A, Mohanty SK. Disease-specific out-of-pocket and catastrophic health expenditure on hospitalization in India: Do Indian households face distress health financing?
- Kumar A, Nagpal J, Bhartia A. Direct cost of ambulatory care of type 2 diabetes in the middle and high income group populace of Delhi: The DEDICOM survey. J Assoc Physicians India.
- Acharya LD, Rau NR, Udupa N, Rajan MS, Vijayanarayana K. Assessment of cost of illness for diabetic patients in South Indian tertiary care hospital. J Pharm Bioallied Sciences.
- Acharya LD, Rau NR, Udupa N, Rajan MS, Vijayanarayana K. Assessment of cost of illness for diabetic patients in South Indian tertiary care hospital. J Pharm Bioallied Sci.
- Rajeshwari S, Adikhari P, Pai MR. Drug utilisation study in geriatric type 2 diabetic patients. J Clin Diagn

A financial candlestick chart on a blue grid. A white curved line, possibly a trend line or moving average, curves across the chart. Several price levels are highlighted with yellow boxes: 104.19, 86.72, and 72.48. A label '61.6 %' is also visible near the top of the curve.

ONLINE COURSE UDEMY.

-
- Statistics for data science and business analytics by 365 carrers.



CONTENTS

- Understand the fundamentals of statistics
- Learn how to work with different types of data
- How to plot different types of data
- Calculate the measures of central tendency, asymmetry, and variability
- Calculate correlation and covariance



-
- Distinguish and work with different types of distributions
 - Estimate confidence intervals
 - Perform hypothesis testing
 - Make data driven decisions
 - Understand the mechanics of regression analysis
 - Carry out regression analysis
 - Use and understand dummy variables
 - Understand the concepts needed for data science even with Python and R!



Learnings.

- How to choose a sample from population data .
- Fundamentals of descriptive statistics.
- Measure of central tendency , asymmetry and variance.
- Central limit theorem
- Distributions .
- Estimators
- Hypothesis testing and its practical applications .



Learnings.

- Regression analysis .
- Corelation and causation.
- Linear regression model.
- Assumptions of linear regression analysis.
- Dealing with categorical data (Dummy variables) and its practical application.

THANK YOU