



Institute of Health Management Research

MBA Hospital and Health Management

Batch-22nd (2017-2019)

First Year

Bio-Statistics

Term Examination



Time - 3 Hour

MM-70

The question paper contains 4 printed pages.

Note:

- a. Kindly attach the question paper with the answer sheet at the time of submission of answer sheet.
- b. All sections are compulsory.
- c. Marks are indicated against each question.

Section A

Kindly tick (✓) against the correct option. (4 x 2marks = 8 marks)

i. Most commonly used statistical average is -

- a. Mean
- b. Median
- c. Mode
- d. None

ii. Mean value of weight of a group of 10 boys was found to be 18.2 kg. Later it was found that weight of one of the boys was wrongly recorded as 2 kg that should have been 20 kg. The true mean weight of the group is -

- a. 18.2 kg
- b. 20.2 kg
- c. 16.4 kg
- d. 20 kg

iii. Systolic BP of a group of persons follows normal distribution curve. The mean BP is 120. The values above 120 are -

- a. 25 per cent
- b. 75 per cent
- c. 50 per cent
- d. 100 per cent

- iv. In a 3×4 contingency table, the number of degrees of freedom equals to -
- 1
 - 5
 - 6
 - 12



Kindly fill in the blanks for the following (3 x 2marks = 6 marks)

- v. Correlation coefficient tends to lie between.....
- vi. If the grading of diabetes is classified as "mild", "moderate" and "severe" the scale of measurement used is.....
- vii. For large degrees of freedom, the distribution becomes almost adistribution

Kindly write True or False against the statement (3 x 2marks = 6 marks)

- viii. The best method to show the association between height and weight of children in a class is by Line diagram
- ix. With increasing degree of accuracy, the sample size increases
- x. Association between an employee's performance level and his job satisfaction could be tested by doing t-test

Section B

Define any four from the following (4 x 5 marks= 20 marks)

- Use of Biostatistics in health and hospital fields
- Levels of measurement (Types of scale)
- Chi squared distribution
- Two types of errors in hypothesis testing
- Correlation
- Steps followed in testing of hypothesis



Section C

Attempt any three from the following questions

1. Suppose the Acme Drug Company develops a new drug, designed to prevent colds. The company states that the drug is equally effective for men and women. To test this claim, they choose a simple random sample of 100 women and 200 men from a population of 100,000 volunteers. At the end of the study, 38% of the women caught a cold; and 51% of the men caught a cold. Based on these findings, can we reject the company's claim that the drug is equally effective for men and women? Use a 0.05 level of significance. (10 marks)
2. Systolic blood pressure (B. P.) of a sample of 566 adult males was taken and it is normally distributed. Mean B. P. was found to be 128.8 mmHg and SD 13.05 mmHg. Construct 95% and 99% confidence interval estimate for the population mean. (10 marks)
3. Systolic blood pressure of 9 normal individuals who had been recumbent for 5 minutes was taken. Then 2 ml of 0.5 per cent solution of hypotensive drug was given and blood pressure recorded again. Did the injection of drug lower the blood pressure? (10 marks)

Serial No.	BP before injection	BP after injection
1.	122	120
2.	121	118
3.	120	115
4.	115	110
5.	126	122
6.	130	130
7.	120	116
8.	125	124
9.	128	125

4. A study was conducted to test if there is any difference between the total energy expenditure in the obese women than that of lean women. Data was collected from 13 lean women and 9 obese women and it is given in the following table –

Table: 24-hour total energy expenditure (MJ/day) in groups of lean and obese women (Prentice et al., 1986)

Descriptive statistics	Lean (n=13)	Obese (n=9)
	6.13	8.79
	7.05	9.19
	7.48	9.21
	7.48	9.68
	7.53	9.69
	7.58	9.97
	7.90	11.51
	8.08	11.85
	8.09	12.79
	8.11	
	8.40	
	10.15	
	10.88	
Mean	8.066	10.298
SD	1.238	1.398

Use appropriate test to answer the question at $\alpha = 0.05$

(10 marks)

5. Five tablets were weighed and then assayed with following results:

Weight (mg)	205	200	202	198	197
Potency (mg)	103	100	101	98	98


- a. Find the regression equation of potency on weight.

(7 marks)

- b. Predict the potency for a 200-mg tablet.

(3 marks)



[illegible]

Standard Normal Probabilities




Table entry for z is the area under the standard normal curve to the left of z .

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

TABLE T/5

Critical Values of χ^2


df	$p = .100$	.050	.025	.010	.005	.001
1	2.71	3.84	5.02	6.63	7.88	10.83
2	4.61	5.99	7.38	9.21	10.60	13.82
3	6.25	7.81	9.35	11.34	12.84	16.27
4	7.78	9.49	11.14	13.28	14.86	18.47
5	9.24	11.07	12.83	15.09	16.75	20.52
6	10.64	12.59	14.45	16.81	18.55	22.46
7	12.02	14.07	16.01	18.48	20.28	24.32
8	13.36	15.51	17.53	20.09	21.96	26.13
9	14.68	16.92	19.02	21.67	23.59	27.88
10	15.99	18.31	20.48	23.21	25.19	29.59
11	17.28	19.68	21.92	24.73	26.76	31.26
12	18.55	21.03	23.34	26.22	28.30	32.91
13	19.81	22.36	24.74	27.69	29.82	34.53
14	21.06	23.68	26.12	29.14	31.32	36.12
15	22.31	25.00	27.49	30.58	32.80	37.70
16	23.54	26.30	28.85	32.00	34.27	39.25
17	24.77	27.59	30.19	33.41	35.72	40.79
18	25.99	28.87	31.53	34.81	37.16	42.31
19	27.20	30.14	32.85	36.19	38.58	43.82
20	28.41	31.41	34.17	37.57	40.00	45.32
21	29.62	32.67	35.48	38.93	41.40	46.80
22	30.81	33.92	36.78	40.29	42.80	48.27
23	32.01	35.17	38.08	41.64	44.18	49.73
24	33.20	36.42	39.36	42.98	45.56	51.18
25	34.38	37.65	40.65	44.31	46.93	52.62
26	35.56	38.89	41.92	45.64	48.29	54.05
27	36.74	40.11	43.19	46.96	49.64	55.48
28	37.92	41.34	44.46	48.28	50.99	56.89
29	39.09	42.56	45.72	49.59	52.34	58.30
30	40.26	43.77	46.98	50.89	53.67	59.70
40	51.81	55.76	59.34	63.69	66.77	73.40
50	63.17	67.50	71.42	76.15	79.49	86.66
60	74.40	79.08	83.30	88.38	91.95	99.61
70	85.53	90.53	95.02	100.43	104.21	112.32
80	96.58	101.88	106.63	112.33	116.32	124.84
90	107.57	113.15	118.14	124.12	128.30	137.21
100	118.50	123.34	129.56	135.81	140.17	149.45

Reprinted with permission from Pearson ES and Hartley HO: Biometrika tables for statisticians, ed 3, 1966.

TABLE T/2

Critical Values of Students' t


df	$p = .1$	.05	.02	.01	.005	.002	.001
1	6.314	12.706	31.821	63.657	127.320	318.31	636.62
2	2.920	4.303	6.965	9.925	14.089	22.327	31.598
3	2.353	3.182	4.541	5.841	7.453	10.214	12.924
4	2.132	2.776	3.747	4.604	5.598	7.173	8.610
5	2.015	2.571	3.365	4.032	4.773	5.893	6.869
6	1.943	2.447	3.143	3.707	4.317	5.208	5.959
7	1.895	2.365	2.998	3.499	4.029	4.785	5.408
8	1.860	2.306	2.896	3.355	3.833	4.501	5.041
9	1.833	2.262	2.821	3.250	3.690	4.297	4.781
10	1.812	2.228	2.764	3.169	3.581	4.144	4.587
11	1.796	2.201	2.718	3.106	3.497	4.025	4.437
12	1.782	2.179	2.681	3.055	3.428	3.930	4.318
13	1.771	2.160	2.650	3.012	3.372	3.852	4.221
14	1.761	2.145	2.624	2.977	3.326	3.787	4.140
15	1.753	2.131	2.602	2.947	3.286	3.733	4.073
16	1.746	2.120	2.583	2.921	3.252	3.686	4.015
17	1.740	2.110	2.567	2.898	3.222	3.646	3.965
18	1.734	2.101	2.552	2.878	3.197	3.610	3.922
19	1.729	2.093	2.539	2.861	3.174	3.579	3.883
20	1.725	2.086	2.528	2.845	3.153	3.552	3.850
21	1.721	2.080	2.518	2.831	3.135	3.527	3.819
22	1.717	2.074	2.508	2.819	3.119	3.505	3.792
23	1.714	2.069	2.500	2.807	3.104	3.485	3.767
24	1.711	2.064	2.492	2.797	3.091	3.467	3.745
25	1.708	2.060	2.485	2.787	3.078	3.450	3.725
26	1.706	2.056	2.479	2.779	3.067	3.435	3.707
27	1.703	2.052	2.473	2.771	3.057	3.421	3.690
28	1.701	2.048	2.467	2.763	3.047	3.408	3.674
29	1.699	2.045	2.462	2.756	3.038	3.396	3.659
30	1.697	2.042	2.457	2.750	3.030	3.385	3.646
40	1.684	2.021	2.423	2.704	2.971	3.307	3.551
60	1.671	2.000	2.390	2.660	2.915	3.232	3.460
120	1.658	1.980	2.358	2.617	2.860	3.160	3.373
∞	1.645	1.960	2.326	2.576	2.807	3.090	3.291

Reprinted with permission from Pearson ES and Hartley HO: Biometrika tables for statisticians, ed 3, 1966.